arXiv:1208.5241v2 [math.AP] 3 Oct 2012

Note on Decaying Turbulence in a Generalised Burgers Equation.

Alexandre Boritchev

Centre de Mathematiques Laurent Schwartz, Ecole Polytechnique, Route de Saclay 91128
Palaiseau Cedex, France.
E-mail: boritchev@math.polytechnique.fr

(Received October 4, 2012)

Abstract. We consider a generalised Burgers equation

$$\frac{\partial u}{\partial t} + f'(u)\frac{\partial u}{\partial x} - \nu \frac{\partial^2 u}{\partial x^2} = 0,\ t \geq 0,\ x \in S^1,$$

where f is strongly convex and ν is small and positive.

Under mild assumptions on the initial condition, we obtain sharp estimates for small-scale quantities. In particular, upper and lower bounds differ only by a multiplicative constant. The quantities which we estimate are the dissipation length scale and averages for structure functions and the energy spectrum for solutions u, which characterise the "Burgulence". Our proof uses a quantitative version of arguments contained in Aurell, Frisch, Lutsko & Vergassola (1992). Our estimates remain true for the inviscid Burgers equation.

Contents

1. Introduction 1
2. Notation and Setup 4
    2.1. Sobolev Spaces 4
    2.2. Preliminaries 5
3. Formulation of the Main Results 5
4. Preliminary estimates 6
5. Estimates for Small-Scale Quantities 10
6. Estimates for Small-Scale Quantities in the Inviscid Limit 16

1. Introduction

The Burgers equation

$$\frac{\partial u}{\partial t} + u\frac{\partial u}{\partial x} - \nu \frac{\partial^2 u}{\partial x^2} = 0, \tag{1.1}$$

where $\nu > 0$ is a constant, appears in many fields of physics and other branches of sciences: see the review articles Bec & Frisch (2001); Bec & Khanin (2007) and references therein; see also Florin (1948).

More importantly for us, this equation is, in some way, the most natural one-dimensional model for the Navier-Stokes system, since both equations have similar nonlinearities and dissipative terms. Despite the fact that the equation (1.1) can be integrated explicitly using the Hopf-Cole transformation, for $\nu \ll 1$, its solutions display non-trivial small-scale behaviour, often referred to as Burgers turbulence or "Burgulence" (Burgers 1974; Chorin 1975; Kida 1979). The language of Kolmogorov 1941 theory (Kolmogorov



FIGURE 1. "Typical" solution of the Burgers equation

1941a,b,c) is traditionally used to describe this behaviour.

Heuristically, it is clear that those solutions display smooth ramps and sharp cliffs (Bec & Frisch 2001; Bec & Khanin 2007), which in the limit $\nu \to 0$ correspond to the N-wave behaviour: see for instance Evans (2008). This is a clear manifestation of small-scale intermittency in space. More precisely, we have the following features for a solution u taken at fixed time t:

- Amplitude of the solution: ~ 1.

- Number of cliffs: ~ 1.

- "Height" of a cliff: ~ -1.

- "Width" of a cliff: $\sim \nu$.

After calculating the Fourier transform of an N-wave, it is natural to conjecture that for a certain range of k energy-type quantities $\frac{1}{2}|\hat{u}_k|^2$ behave, in average, as k^{-2} (Chorin 1975; Fournier & Frisch 1983; Kida 1979; Kraichnan 1968).

In the physical space, the natural analogues of small-scale quantities $\frac{1}{2}|\hat{u}_k|^2$ are the structure functions

$$S_p(\ell) = \int_{S^1} (u(x+\ell) - u(x))^p \; dx.$$

For $p \geq 0$, the description above implies that for $\nu \ll \ell \ll 1$, these quantities behave as $\ell^{\max(1,p)}$: in other words, we have a *bifractal behaviour* (Frisch 1995, Chapter 8). Indeed, there are three possibilities for the interval $[x, x+\ell]$:

- $[x, x+\ell]$ covers a large part of a "cliff".

Probability $\simeq C\ell$. $u(x+\ell) - u(x) = \underbrace{-C}_{cliff} + \underbrace{C\ell}_{ramps} = -C$. $|u(x+\ell) - u(x)|^p \overset{p}{\sim} 1.$

- $[x, x+\ell]$ covers a small part of a "cliff".

Contribution of this term is negligible.

- $[x, x+\ell]$ does not intersect a "cliff".

Probability $\simeq 1 - C\ell$. $u(x+\ell) - u(x) = \underbrace{C\ell}_{ramp}$. $|u(x+\ell) - u(x)|^p \overset{p}{\sim} \ell^p$.

Thus, $S_p(\ell) \simeq C(p)(\ell + \ell^p) \overset{p}{\sim} \begin{cases} \ell^p, \ 0 \leq p \leq 1. \\ \ell, \ p \geq 1. \end{cases}$

In this paper, we consider a generalised one-dimensional space-periodic Burgers equation

$$\frac{\partial u}{\partial t} + \frac{df(u)}{dx} - \nu \frac{\partial^2 u}{\partial x^2} = 0, \quad x \in S^1 = \mathbb{R}/\mathbb{Z} \tag{1.2}$$

with f C^∞-smooth and strongly convex, i.e. satisfying

$$f''(y) \geq \sigma > 0, \quad y \in \mathbb{R}. \tag{1.3}$$

The classical Burgers equation (1.1) corresponds to $f(u) = u^2/2$. For simplicity, we only consider solutions with zero space average for fixed t:

$$\int_{S^1} u(t,x)dx = 0, \quad \forall t \geq 0. \tag{1.4}$$

Estimating small-scale quantities for nonlinear PDEs with small viscosity is motivated by the problem of turbulence. This research was initiated by Kuksin, who obtained estimates for a large class of equations (see Kuksin (1997, 1999) and references in Kuksin (1999)). Biryuk (2001) obtained some estimates for solutions of (1.2) which naturally depend on the initial data $u_0 = u(0, \cdot)$, since this equation is dissipative for the energy $\int_{S^1} u^2/2$ because of the damping term $\nu \partial^2 u / \partial x^2$. In Boritchev (2012*a*,*b*), we obtain similar results which are independent of u_0. However, in both articles we add a rough in time and smooth in space random forcing term in the right-hand side of (1.2) (a "kicked" and a white force, respectively). Thus, we change the nature of the equation: the energy injection due to the random forcing now balances the dissipation. Note also that in E, Khanin, Mazel & Sinai (1997), the authors obtain similar results for the stationary solution in the limit $\nu \to 0$, for a forcing of the same type as in Boritchev (2012*b*).

Our estimates, which confirm the predictions for values of small-scale qualities listed above, are very similar to the ones in Biryuk (2001) and Boritchev (2012*a*,*b*) and depend on the initial data through the constant

$$D = \max(|u_0|_1^{-1}, \ |u_0|_{1,\infty}).$$

The physical meaning of the quantity D is that it yields an upper bound for the ratio between the maximal rate of dissipation and the amount of energy initially contained in the system.

After introducing the notation and setup in Section 2, we formulate the main results in Section 3. In Section 4, we begin by estimating from above the quantity $\partial u / \partial x$. This crucial bound allows us to obtain upper bounds, as well as time-averaged lower bounds, of Sobolev norms $|u|_{m,p}$. Those bounds depend only on f and the constant D.

In Section 5, we give sharp upper and lower bounds for dissipation length scale, structure functions, flatness, and energy spectrum for the flow $u(t,x)$, which hold uniformly for $\nu \leq \nu_0$, and we analyse the meaning of these results in terms of the theory of turbulence. Those bounds and the constant $\nu_0 > 0$ only depend on f and D.

These results rigorously justify the previously stated classical predictions for small-scale statistical quantities for decaying Burgers turbulence. In Section 6, we show that the results in Section 5 still hold in the inviscid limit $\nu \to 0$. Many proofs in this paper are very similar to the ones in Biryuk (2001) or Boritchev (2012*b*). Accordingly sometimes we omit the proof and instead refer the reader to those two papers.

Note that when studying the typical behaviour of PDEs, one usually considers some averaging in the initial condition. The idea is to avoid pathological cases, since there is no random mechanism to get solutions out of "bad" regions. Here, no such averaging is actually necessary. This is due to the particular structure of the deterministic Burgers equation: an initial condition u_0 is as "generic" as the ratio between the orders of $(u_0)_x$ and of u_0 itself, which can be bounded from above using the quantity D.

2. Notation and Setup

Agreement: In the whole paper, all functions that we consider are real-valued and, since we work on $S^1 = \mathbb{R}/\mathbb{Z}$, periodic.

2.1. *Sobolev Spaces*

Consider a zero mean value integrable function v on S^1. For $p \in [1, \infty)$, we denote its L_p norm

$$\Big(\int_{S^1} |v|^p \Big)^{1/p}$$

by $|v|_p$. The L_∞ norm is by definition

$$|v|_\infty = \max_{x \in S^1} |v(x)|.$$

The L_2 norm is denoted by $|v|$, and $\langle \cdot, \cdot \rangle$ stands for the L_2 scalar product. From now on $L_p,\ p \in [1, \infty]$ denotes the space of zero mean value functions in $L_p(S^1)$. Similarly, C^∞ is the space of C^∞-smooth zero mean value functions on S^1.

For a nonnegative integer m and $p \in [1, \infty]$, $W^{m,p}$ stands for the Sobolev space of zero mean value functions v on S^1 with finite norm

$$|v|_{m,p} = \left| \frac{d^m v}{dx^m} \right|_p .$$

In particular, $W^{0,p} = L_p$ for $p \in [1, \infty]$. For $p = 2$, we denote $W^{m,2}$ by H^m, and abbreviate the corresponding norm as $\|v\|_m$.

Note that since the length of S^1 is 1 and the mean value of v vanishes, we have

$$|v|_1 \leq |v|_\infty \leq |v|_{1,1} \leq |v|_{1,\infty} \leq \cdots \leq |v|_{m,1} \leq |v|_{m,\infty} \leq \ldots$$

We recall a version of the classical Gagliardo-Nirenberg inequality: cf. Doering & Gibbon (1995, Appendix).

Lemma 2.1. *For a smooth zero mean value function v on S^1,*

$$|v|_{\beta,r} \leq C\, |v|^{\theta}_{m,p}\, |v|^{1-\theta}_q ,$$

where $m > \beta$, and r is defined by

$$\frac{1}{r} = \beta - \theta\Big(m - \frac{1}{p}\Big) + (1-\theta)\frac{1}{q},$$

under the assumption $\theta = \beta/m$ if $p = 1$ or $p = \infty$, and $\beta/m \leq \theta < 1$ otherwise. The constant C depends on m, p, q, β, θ.

Subindices t and x, which can be repeated, denote partial differentiation with respect to the corresponding variables. We denote by $v^{(m)}$ the m-th derivative of v in the variable x. The function $v(t, \cdot)$ is abbreviated as $v(t)$.

2.2. *Preliminaries*

In this paper, we study asymptotical properties of solutions to (1.2) for small values of ν, i.e. we suppose that

$$0 \ll \nu \leq 1.$$

We assume that f is infinitely derivable and satisfies (1.3). We recall that we restrict ourselves to the zero space average case, i.e. the initial condition $u_0 := u(0,\cdot)$ satisfies (1.4). Consequently, $u(t)$ satisfies (1.4) for all t. Finally, we assume that $u_0 \in C^\infty$, and we denote by D the quantity

$$D = \max(|u_0|_1^{-1},\ |u_0|_{1,\infty}). \tag{2.1}$$

In particular, we assume that we are not in the case $u_0 \equiv 0$, corresponding to the trivial solution $u(t,x) \equiv 0$. Note that $D > 1$. Moreover, for $0 \leq m \leq 1$ and $1 \leq p \leq \infty$, we have

$$D^{-1} \leq |u_0|_{m,p} \leq D. \tag{2.2}$$

For the existence, uniqueness, and smoothness of solutions to (1.2), see for instance Kreiss & Lorenz (1989).

Agreements: From now on, all constants denoted by C with sub- or superindexes are strictly positive. Unless otherwise stated, they depend only on f and the constant D. By $C(a_1,\dots,a_k)$ we denote constants which also depend on parameters $a_1,\dots,a_k$. By $X \overset{a_1,\dots,a_k}{\lesssim} Y$ we mean that $X \leq C(a_1,\dots,a_k)Y$. The notation $X \overset{a_1,\dots,a_k}{\sim} Y$ stands for

$$Y \overset{a_1,\dots,a_k}{\lesssim} X \overset{a_1,\dots,a_k}{\lesssim} Y.$$

In particular, $X \lesssim Y$ and $X \sim Y$ mean that $X \leq CY$ and $C^{-1}Y \leq X \leq CY$, respectively.

All constants are independent of the viscosity ν. We denote by $u = u(t,x)$ a solution of (1.2) for an initial condition u_0. A relation where the admissible values of t (respectively, x) are not specified is assumed to hold for all $t \geq 0$ or $t > 0$, depending on the context (respectively, all $x \in S^1$).

The brackets $\{\cdot\}$ stand for averaging in time over an interval $[T_1, T_2]$, where T_1, T_2 only depend on f and D (see (4.11) for their definition.)

We use the notation $g^- = \max(-g, 0)$ and $g^+ = \max(g, 0)$.

3. Formulation of the Main Results

In Section 4, we prove sharp upper and lower bounds for some Sobolev norms of u. The key estimate is obtained in Lemma 4.1: we show there that

$$u_x(t,x) \leq \min(D, \sigma^{-1}t^{-1}). \tag{3.1}$$

The main results for Sobolev norms of solutions are summed up in Theorem 4.7. Namely, for $m \geq 0$ we have

$$\Big(\{|u(t)|^\alpha_{m,\infty}\}\Big)^{1/\alpha} \overset{m,\alpha}{\sim} \nu^{-m}, \quad \alpha > 0, \tag{3.2}$$

where $\{\cdot\}$ denotes averaging in time over the interval $[T_1, T_2]$ defined by (4.11).

In Section 5 we obtain sharp estimates for analogues of the quantities characterising the hydrodynamical turbulence. In what follows, we assume that $\nu \in (0, \nu_0]$, where $\nu_0 \in (0,1]$ only depends on f and D.

To begin with, we define the non-empty and non-intersecting intervals

$$J_1 = (0,\ C_1\nu];\ J_2 = (C_1\nu,\ C_2];\ J_3 = (C_2,\ 1]$$



Figure 2. Scales for the 1D Burgers solutions

corresponding to the *dissipation range*, the *inertial range*, and the *energy range* from the Kolmogorov 1941 theory of turbulence, respectively: see Frisch (1995). Then we consider the averaged moments of increments in space for the flow $u(t,x)$:

$$S_p(\ell) = \left\{ \int_{S^1} |u(t,x+\ell) - u(t,x)|^p dx \right\},\ 0 < \ell \leq 1.$$

The quantity $S_p(\ell)$ is (up to averaging) the *structure function* of p-th order. As a consequence of (3.1-3.2), in Theorem 5.9 we prove that for $\ell \in J_1$:

$$S_p(\ell) \overset{p}{\sim} \begin{cases} \ell^p,\ 0 \leq p \leq 1. \\ \ell^p \nu^{-(p-1)},\ p \geq 1, \end{cases}$$

and for $\ell \in J_2$:

$$S_p(\ell) \overset{p}{\sim} \begin{cases} \ell^p,\ 0 \leq p \leq 1. \\ \ell,\ p \geq 1. \end{cases}$$

Consequently, for $\ell \in J_2$ the flatness satisfies:

$$F(\ell) := S_4(\ell)/S_2^2(\ell) \sim \ell^{-1}.$$

Thus, u is highly intermittent in the inertial range. This intermittency is due to the particular structure of the solution, where the excited zones correspond to the cliffs: cf. Frisch (1995) for a discussion of intermittency for hydrodynamical turbulent flows.

Finally, we get estimates for the spectral asymptotics of Burgulence. On one hand, as a consequence of Lemma 4.3, for $m \geq 1$ we get:

$$\{|\hat{u}_k|^2\} \overset{m}{\lesssim} k^{-2m} \|u\|_m^2 \overset{m}{\lesssim} (k\nu)^{-2m}\nu.$$

In particular, $\{|\hat{u}_k|^2\}$ decreases at a faster-than-algebraic rate for $|k| \succeq \nu^{-1}$. On the other hand, by Theorem 5.11 for k such that $k^{-1} \in J_2$ the *energy spectrum* $E(k)$ satisfies

$$E(k) = \left\{ \frac{\sum_{|n| \in [M^{-1}k, Mk]} |\hat{u}_n|^2}{\sum_{|n| \in [M^{-1}k, Mk]} 1} \right\} \sim k^{-2},$$

where $M \geq 1$ is a constant depending only on f and D.

Finally, we note that estimates for small-scale quantities still hold in the inviscid limit $\nu \to 0$, up to some natural modifications (see Section 6).

4. Preliminary estimates

We begin by proving a key upper estimate for u_x.

Lemma 4.1. *We have*

$$u_x(t,x) \leq \min(D, \sigma^{-1} t^{-1}).$$

Proof. Differentiating the equation (1.2) once in space we get

$$(u_x)_t + f''(u)u_x^2 + f'(u)(u_x)_x = \nu(u_x)_{xx}.$$

Now consider a strictly positive point of maximum (t_1, x_1) for u_x on the cylinder $S = [0,t] \times S^1$, such that $t_1 > 0$. At such a point, we would have $(u_x)_t \geq 0$, $(u_x)_x = 0$, and $(u_x)_{xx} \leq 0$. Consequently, by (1.3) we get

$$\sigma u_x^2 \leq f''(u)u_x^2 \leq 0,$$

which is impossible. Thus u_x can only reach a strictly positive maximum on S for $t_1 = 0$. In other words, we have

$$u_x(t,x) \leq \max_{x \in S^1} (u_0)_x(x) \leq D.$$

The inequality

$$u_x(t,x) \leq \sigma^{-1} t^{-1}$$

is proved in by a similar maximum principle argument applied to the function tu_x: cf. Kruzhkov (1964). □

Since the space averages of $u(t)$ and $u_x(t)$ vanish, we get the following upper estimates:

$$|u(t)|_p \leq |u(t)|_\infty \leq \int_{S^1} u_x^+(t) \leq \min(D, \sigma^{-1}t^{-1}), \quad 1 \leq p \leq +\infty, \tag{4.1}$$

$$|u(t)|_{1,1} = 2\int_{S^1} u_x^+(t) \leq 2\min(D, \sigma^{-1}t^{-1}). \tag{4.2}$$

Now we recall a standard estimate for the nonlinearity $\left\langle v^{(m)}, (f(v))^{(m+1)} \right\rangle$. For its proof, we refer to Boritchev (2012*b*).

Lemma 4.2. *For $v \in C^\infty$ such that $|v|_\infty \leq A$, we have*

$$\left| \left\langle v^{(m)}, (f(v))^{(m+1)} \right\rangle \right| \leq \tilde{C} \left\| v \right\|_m \left\| v \right\|_{m+1}, \quad m \geq 1,$$

where $\tilde{C}$ depends only on m, A, and $|f|_{C^m([-A,A])}$.

The following result shows that there is a strong nonlinear damping which prevents the successive derivatives of u from becoming too large.

Lemma 4.3. *We have*

$$\left\| u(t) \right\|_1^2 \lesssim \nu^{-1}.$$

On the other hand, for $m \geq 2$,

$$\left\| u(t) \right\|_m^2 \overset{m}{\lesssim} \max(\nu^{-(2m-1)}, t^{-(2m-1)}).$$

Proof. Fix $m \geq 1$. Denote

$$x(t) = \left\| u(t) \right\|_m^2.$$

We claim that the following implication holds:

$$x(t) \geq C'\nu^{-(2m-1)} \Longrightarrow \frac{d}{dt}x(t) \leq -(2m-1)x(t)^{2m/(2m-1)}, \tag{4.3}$$

where C' is a fixed strictly positive number, chosen later. Below, all constants denoted by C do not depend on C'.

Indeed, assume that $x(t) \geq C'\nu^{-(2m-1)}$. Integrating by parts in space and using (4.1) ($p=\infty$) and Lemma 4.2, we get the following energy dissipation relation:

$$\begin{aligned}\frac{d}{dt}x(t) &= -2\nu \left\|u(t)\right\|_{m+1}^2 - 2\left\langle u^{(m)}(t), (f(u(t)))^{(m+1)}\right\rangle \\ &\leq -2\nu \left\|u(t)\right\|_{m+1}^2 + C\left\|u(t)\right\|_m \left\|u(t)\right\|_{m+1}. \end{aligned} \tag{4.4}$$

Applying Lemma 2.1 to u_x and then using (4.2), we get

$$\begin{aligned}\left\|u(t)\right\|_m &\leq C\left\|u(t)\right\|_{m+1}^{(2m-1)/(2m+1)} |u(t)|_{1,1}^{2/(2m+1)} \\ &\leq C\left\|u(t)\right\|_{m+1}^{(2m-1)/(2m+1)}. \end{aligned} \tag{4.5}$$

Thus, we have the relation

$$\frac{d}{dt}x(t) \leq (-2\nu \left\|u(t)\right\|_{m+1}^{2/(2m+1)} + C)\left\|u(t)\right\|_{m+1}^{4m/(2m+1)}. \tag{4.6}$$

The inequality (4.5) yields

$$\left\|u(t)\right\|_{m+1}^{2/(2m+1)} \geq Cx(t)^{1/(2m-1)}, \tag{4.7}$$

and then since by assumption $x(t) \geq C'\nu^{-(2m-1)}$ we get

$$\left\|u(t)\right\|_{m+1}^{2/(2m+1)} \geq CC'^{1/(2m-1)}\nu^{-1}. \tag{4.8}$$

Combining the inequalities (4.6-4.8), for C' large enough we get

$$\frac{d}{dt}x(t) \leq (-CC'^{1/(2m-1)} + C)x(t)^{2m/(2m-1)}.$$

Thus we can choose C' in such a way that the implication (4.3) holds.

For $m=1$, (2.2) and (4.3) immediately yield that

$$x(t) \leq \max(C'\nu^{-1}, D^2) \leq \max(C', D^2)\nu^{-1},\ t \geq 0.$$

Now consider the case $m \geq 2$. We claim that

$$x(t) \leq \max(C'\nu^{-(2m-1)}, t^{-(2m-1)}). \tag{4.9}$$

Indeed, if $x(s) \leq C'\nu^{-(2m-1)}$ for some $s \in [0,t]$, then the assertion (4.3) ensures that $x(s)$ remains below this threshold up to time t.

Now, assume that $x(s) > C'\nu^{-(2m-1)}$ for all $s \in [0,t]$. Denote

$$\tilde{x}(s) = (x(s))^{-1/(2m-1)},\ s \in [0,t].$$

By (4.3) we get $d\tilde{x}(s)/ds \geq 1$. Therefore $\tilde{x}(t) \geq t$, and $x(t) \leq t^{-(2m-1)}$. Thus in this case, inequality (4.9) still holds. This proves the lemma's assertion. □

By Lemma 2.1 applied to $u^{(m)}$ we get the following inequality for $m \geq 1$:

$$|u(t)|_{m,\infty} \lesssim \left\|u(t)\right\|_m^{1/2} \left\|u(t)\right\|_{m+1}^{1/2} \overset{m}{\lesssim} \max(\nu^{-m}, t^{-m}). \tag{4.10}$$

Now we define

$$T_1 = \frac{1}{4}D^{-2}\tilde{C}^{-1}; \quad T_2 = \max\Big(\frac{3}{2}T_1, \quad 2D\sigma^{-1}\Big), \tag{4.11}$$

where $\tilde{C}$ is a constant such that for all t, $\left\|u(t)\right\|_1^2 \leq \tilde{C}\nu^{-1}$ (see Lemma 4.3). From now on, for any function $A(t)$, $\{A(t)\}$ is by definition the time average

$$\{A(t)\} = \frac{1}{T_2 - T_1}\int_{T_1}^{T_2} A(t).$$

The first quantity that we estimate from below is $\{\|u(t)\|_1^2\}$.

LEMMA 4.4. *We have*

$$\{\|u(t)\|_1^2\} \gtrsim \nu^{-1}.$$

Proof. Integrating by parts in space, we get the dissipation identity

$$\frac{d}{dt}|u(t)|^2 = \int_{S^1} (-2uf'(u)u_x + 2\nu u u_{xx}) = -2\nu \|u(t)\|_1^2. \tag{4.12}$$

Thus, integrating in time and using (2.1) and Lemma 4.3, we obtain that

$$|u(T_1)|^2 = |u_0|^2 - 2\nu \int_0^{T_1} \|u(t)\|_1^2 \geq D^{-2} - 2T_1\tilde{C} \geq \frac{1}{2}D^{-2}.$$

Consequently, integrating (4.12) in time and using (4.1) ($p = 2$) we get

$$\begin{aligned}
\{\|u(t)\|_1^2\} &= \frac{1}{2\nu(T_2 - T_1)}(|u(T_1)|^2 - |u(T_2)|^2) \\
&\geq \frac{1}{2\nu(T_2 - T_1)}\Big(\frac{1}{2}D^{-2} - \sigma^{-2}T_2^{-2}\Big) \\
&\geq \frac{D^{-2}}{8(T_2 - T_1)}\nu^{-1},
\end{aligned}$$

which proves the lemma's assertion. □

This time-averaged lower bound yields similar bounds for other Sobolev norms.

LEMMA 4.5. *For* $m \geq 1$,

$$\{\|u(t)\|_m^2\} \overset{m}{\gtrsim} \nu^{-(2m-1)}.$$

Proof. Since the case $m = 1$ has been treated in the previous lemma, we may assume that $m \geq 2$. By (4.2) and Lemma 2.1, we have:

$$\{\|u(t)\|_m^2\} \overset{m}{\gtrsim} \{\|u(t)\|_m^2\, |u(t)|_{1,1}^{(4m-4)}\} \overset{m}{\gtrsim} \{\|u(t)\|_1^{4m-2}\}.$$

Thus, using Hölder's inequality and Lemma 4.4, we get:

$$\{\|u(t)\|_m^2\} \overset{m}{\gtrsim} \{\|u(t)\|_1^{4m-2}\} \overset{m}{\gtrsim} \{\|u(t)\|_1^2\}^{(2m-1)} \overset{m}{\gtrsim} \nu^{-(2m-1)}. \ \square$$

The following lemma is proved similarly.

LEMMA 4.6. *For* $m \geq 0$, *we have:*

$$\{|u(t)|_{m,\infty}^2\} \overset{m}{\gtrsim} \nu^{-2m}.$$

The following theorem sums up the results of this section which will be used later, with the exception of Lemma 4.1.

THEOREM 4.7. *For* $m \geq 0$, *we have:*

$$\Big(\{|u(t)|_{m,\infty}^{\alpha}\}\Big)^{1/\alpha} \overset{m,\alpha}{\sim} \nu^{-m}, \qquad \alpha > 0, \tag{4.13}$$

where $\{\cdot\}$ denotes time-averaging over $[T_1, T_2]$. The upper estimates in (4.13) hold without time-averaging, uniformly for t separated from 0. Namely, we have

$$|u(t)|_{m,\infty} \overset{m}{\lesssim} \max(\nu^{-m}, t^{-m}).$$

On the other hand, the lower estimates hold for all m, α.

Proof. Upper estimates follow from the inequality (4.10). Lower estimates for $\alpha \geq 2$ follow from Lemma 4.6 by Hölder's inequality. Finally, we obtain lower estimates for $\alpha \in (0,2)$ using lower estimates for $\alpha = 2$, upper estimates for $\alpha = 3$, and Hölder's inequality. Indeed:

$$\{|u(t)|^{\alpha}_{m,\infty}\} \geq \Big(\{|u(t)|^{2}_{m,\infty}\}\Big)^{3-\alpha} \Big(\{|u(t)|^{3}_{m,\infty}\}\Big)^{-(2-\alpha)}. \ \square$$

5. Estimates for Small-Scale Quantities

In this section, we study analogues of quantities which are important for the study of hydrodynamical turbulence. We consider quantities in physical space (structure functions, flatness) as well as in Fourier space (energy spectrum). We assume that $\nu \leq \nu_0$. The value of $\nu_0 > 0$ will be chosen in (5.10).

We define the intervals

$$J_1 = (0,\ C_1\nu];\ J_2 = (C_1\nu,\ C_2];\ J_3 = (C_2,\ 1].$$

The strictly positive constants C_1 and C_2 will be chosen in (5.9-5.10) in such a manner that $C_1\nu_0 < C_2 < 1$, which ensures that J_i are non-empty and non-intersecting.

We observe that $\{|u|^2\} \gtrsim 1$, since $|u|^2 \geq D^{-2}/4$ on $[T_1, 3T_1/2]$ (see proof of Lemma 4.4). Thus, by Theorem 4.7, we obtain that $\{|u|^2\} \sim 1$. On the other hand, by (4.2) (after integration by parts) we get

$$\{|\hat{u}_n|^2\} = (2\pi n)^{-2} \Big\{ \Big| \int_{S^1} e^{2\pi i n x} u_x(x) \Big|^2 \Big\} \leq (2\pi n)^{-2} \{|u|^2_{1,1}\} \leq C n^{-2}, \qquad (5.1)$$

and C_1 and C_2 can be made as small as desired (cf. (5.11)). Consequently, the proportion of the sum $\{\sum |\hat{u}_n|^2\}$ contained in Fourier modes corresponding to J_3 can be made as large as desired. For instance, we may assume that

$$\Bigg\{ \sum_{|n| \leq C_2^{-1}} |\hat{u}_n|^2 \Bigg\} \geq \frac{99}{100} \Bigg\{ \sum_{n \in \mathbb{Z}} |\hat{u}_n|^2 \Bigg\}.$$

For $p \geq 0$, we set:

$$S_p(\ell) = \Big\{ \int_{S^1} |u(t, x+\ell) - u(t,x)|^p dx \Big\}.$$

This quantity corresponds (up to averaging) to the structure function of p-th order. The flatness $F(\ell)$, which measures spatial intermittency, is given by

$$F(\ell) = S_4(\ell)/S_2^2(\ell), \qquad (5.2)$$

see Frisch (1995). Finally, for $k \geq 1$, we define the (layer-averaged) energy spectrum by

$$E(k) = \Bigg\{ \frac{\sum_{|n| \in [M^{-1}k, Mk]} |\hat{u}_n|^2}{\sum_{|n| \in [M^{-1}k, Mk]} 1} \Bigg\}, \qquad (5.3)$$

where $M \geq 1$ is a constant which will be specified later (see the proof of Theorem 5.11).

We begin by estimating the functions $S_p(\ell)$ from above.

Lemma 5.1. *For* $\ell \in [0,1]$,

$$S_p(\ell) \overset{p}{\lesssim} \begin{cases} \ell^p, \ 0 \leq p \leq 1. \\ \ell^p \nu^{-(p-1)}, \ p \geq 1. \end{cases}$$

Proof. We begin by considering the case $p \geq 1$. We have

$$\begin{aligned} S_p(\ell) &= \Big\{ \int_{S^1} |u(x+\ell) - u(x)|^p dx \Big\} \\ &\leq \Big\{ \Big(\int_{S^1} |u(x+\ell) - u(x)| dx \Big) \Big(\max_x |u(x+\ell) - u(x)|^{p-1} \Big) \Big\}. \end{aligned}$$

Using the fact that the space average of $u(x+\ell) - u(x)$ vanishes and Hölder's inequality, we obtain that

$$\begin{aligned} S_p(\ell) \leq & \Big\{ \Big(2 \int_{S^1} (u(x+\ell) - u(x))^+ dx \Big)^p \Big\}^{1/p} \times \Big\{ \max_x |u(x+\ell) - u(x)|^p \Big\}^{(p-1)/p} \\ \leq & C\ell \Big\{ \max_x |u(x+\ell) - u(x)|^p \Big\}^{(p-1)/p}, \end{aligned} \tag{5.4}$$

where the second inequality follows from Lemma 4.1. Finally, by Theorem 4.7 we get

$$S_p(\ell) \leq C\ell \Big\{ (\ell |u|_{1,\infty})^p \Big\}^{(p-1)/p} \leq C\ell^p \nu^{-(p-1)}.$$

The case $p < 1$ follows immediately from the case $p = 1$ since now $S_p(\ell) \leq (S_1(\ell))^p$, by Hölder's inequality. □

For $\ell \in J_2 \cup J_3$, we have a better upper bound if $p \geq 1$.

Lemma 5.2. *For* $\ell \in J_2 \cup J_3$,

$$S_p(\ell) \overset{p}{\lesssim} \begin{cases} \ell^p, \ 0 \leq p \leq 1. \\ \ell, \ p \geq 1. \end{cases}$$

Proof. The calculations are almost the same as in the previous lemma. The only difference is that we use another bound for the right-hand side of (5.4). Namely, by Theorem 4.7 we have

$$\begin{aligned} S_p(\ell) &\leq C\ell \Big\{ \max_x |u(x+\ell) - u(x)|^p \Big\}^{(p-1)/p} \\ &\leq C\ell \Big\{ (2|u|_\infty)^p \Big\}^{(p-1)/p} \leq C\ell. \ \square \end{aligned}$$

Remark 5.3. *The Lemmas 5.1 and 5.2 actually hold even if we drop the time-averaging, since in deriving them we only use upper estimates which hold uniformly for* $t \geq T_1$.

To prove the lower estimates for $S_p(\ell)$, we need a lemma. Loosely speaking, this lemma states that there exists a large enough set $L_K \subset [T_1, T_2]$ such that for $t \in L_K$, several Sobolev norms are of the same order as their time averages. Thus, for $t \in L_K$, we can prove the existence of a "cliff" of height at least C and width at least $C\nu$, using some of the arguments in Aurell, Frisch, Lutsko & Vergassola (1992) which we explained in the introduction.

Note that in the following definition, (5.5-5.6) contain lower and upper estimates, while (5.7) only contains an upper estimate. The inequality $|u(t)|_\infty \leq \max u_x(t)$ in (5.5) always holds, since $u(t)$ has zero mean value and the length of S^1 is 1.

Definition 5.4. *For $K > 1$, we denote by L_K the set of all $t \in [T_1, T_2]$ such that the assumptions*

$$K^{-1} \leq |u(t)|_\infty \leq \max u_x(t) \leq K \tag{5.5}$$

$$K^{-1}\nu^{-1} \leq |u(t)|_{1,\infty} \leq K\nu^{-1} \tag{5.6}$$

$$|u(t)|_{2,\infty} \leq K\nu^{-2} \tag{5.7}$$

hold.

Lemma 5.5. *There exist constants $C, K_1 > 0$ such that for $K \geq K_1$, the Lebesgue measure of L_K verifies $\lambda(L_K) \geq C$.*

Proof. We begin by noting that if $K \leq K'$, then $L_K \subset L_{K'}$. By Lemma 4.1 and Theorem 4.7, for K large enough the upper estimates in (5.5-5.7) hold for all t. Therefore, if we denote by B_K the set of t such that

"The lower estimates in (5.5-5.6) hold for a given value of K",

then it suffices to prove the lemma's statement with B_K in place of L_K. Now denote by D_K the set of t such that

"The lower estimate in (5.6) holds for a given value of K".

By Lemma 2.1 we have

$$|u|_\infty \geq C|u|_{2,\infty}^{-1}|u|_{1,\infty}^2.$$

Thus, if D_K holds, then $B_{K'}$ holds for K' large enough. Now it remains to show that there exists $C > 0$ such that for K large enough, we have the inequality $\lambda(D_K) \geq C$. We clearly have

$$\{|u|_{1,\infty}\mathbf{1}(|u|_{1,\infty} \leq K^{-1}\nu^{-1})\} \leq K^{-1}\nu^{-1}.$$

Here, $\mathbf{1}(A)$ denotes the indicator function of an event A. On the other hand, by the estimate for $\{|u|_{1,\infty}^2\}$ in Theorem 4.7 we get

$$\{|u|_{1,\infty}\mathbf{1}(|u|_{1,\infty} \geq K\nu^{-1})\} \leq K^{-1}\nu\{|u|_{1,\infty}^2\} \leq CK^{-1}\nu^{-1}$$

Now denote by f the function

$$f = |u|_{1,\infty}\mathbf{1}(K_0^{-1}\nu^{-1} \leq |u|_{1,\infty} \leq K_0\nu^{-1}).$$

The inequalities above and the lower estimate for $\{|u|_{1,\infty}\}$ in Theorem 4.7 imply that

$$\{f\} \geq (C - K_0^{-1} - CK_0^{-1})\nu^{-1} \geq C_0\nu^{-1},$$

for some suitable constants C_0 and K_0. Since $f \leq K_0\nu^{-1}$, then we get

$$\lambda(f \geq C_0\nu^{-1}/2) \geq C_0K_0^{-1}(T_2 - T_1)/2.$$

Thus, since $|u|_{1,\infty} \geq f$, we have the inequality

$$\lambda(|u|_{1,\infty} \geq C_0\nu^{-1}/2) \geq C_0K_0^{-1}(T_2 - T_1)/2,$$

which implies existence of $C, K_1 > 0$ such that $\lambda(D_K) \geq C$ for $K \geq K_1$. $\square$

Let us denote by $O_K \subset [T_1, T_2]$ the set defined as L_K, but with relation (5.6) replaced by

$$K^{-1}\nu^{-1} \leq -\min u_x \leq K\nu^{-1}. \tag{5.8}$$

Corollary 5.6. *For $K \geq K_1$ and $\nu < K_1^{-2}$, we have $\lambda(O_K) \geq C$.*

Proof. For $K = K_1$ and $\nu < K_1^{-2}$, the estimates (5.5-5.6) tell us that

$$\max u_x(t) \leq K_1 < K_1^{-1}\nu^{-1} \leq |u_x(t)|_\infty, \quad t \in L_K.$$

Thus, in this case the assertion of Lemma 5.5 with (5.6) replaced by (5.8) holds for the set $O_K = L_K$. Since increasing K while keeping ν constant increases the measure of O_K, for $K \geq K_1$ and $\nu < K_1^{-2}$ we still have $\lambda(O_K) \geq C$. □

Now we fix

$$K = K_1, \tag{5.9}$$

and choose

$$\nu_0 = \frac{1}{6}K^{-2};\ C_1 = \frac{1}{4}K^{-2};\ C_2 = \frac{1}{20}K^{-4}. \tag{5.10}$$

In particular, we have $0 < C_1\nu_0 < C_2 < 1$: thus the intervals J_i are non-empty and non-intersecting for all $\nu \in (0, \nu_0]$. Everywhere below the constants depend on K.

Actually, we can choose any values of C_1, C_2, and ν_0, provided

$$C_1 \leq \frac{1}{4}K^{-2}; \quad 5K^2 \leq \frac{C_1}{C_2} < \frac{1}{\nu_0}. \tag{5.11}$$

Lemma 5.7. *For $\ell \in J_1$,*

$$S_p(\ell) \overset{p}{\gtrsim} \begin{cases} \ell^p,\ 0 \leq p \leq 1. \\ \ell^p \nu^{-(p-1)},\ p \geq 1. \end{cases}$$

Proof. By Corollary 5.6, it suffices to prove that the inequalities hold uniformly in t for $t \in O_K$, with $S_p(\ell)$ replaced by

$$\int_{S^1} |u(x+\ell) - u(x)|^p dx.$$

Till the end of this proof, we assume that $t \in O_K$.

Denote by z the leftmost point on S^1 (considered as $[0,1)$) such that $u'(z) \leq -K^{-1}\nu^{-1}$. Since $|u|_{2,\infty} \leq K\nu^{-2}$, we have

$$u'(y) \leq -\frac{1}{2}K^{-1}\nu^{-1}, \quad y \in [z - \frac{1}{2}K^{-2}\nu, z + \frac{1}{2}K^{-2}\nu]. \tag{5.12}$$

In other words, the interval

$$[z - \frac{1}{2}K^{-2}\nu, z + \frac{1}{2}K^{-2}\nu]$$

corresponds to (a part of) a cliff.

Case $\mathbf{p \geq 1}$. Since $\ell \leq C_1\nu = \frac{1}{4}K^{-2}\nu$, then by Hölder's inequality we get

$$\begin{aligned}
&\int_{S^1} |u(x+\ell)-u(x)|^p dx \geq \int_{z-\frac{1}{4}K^{-2}\nu}^{z+\frac{1}{4}K^{-2}\nu} |u(x+\ell)-u(x)|^p dx \\
&\geq (K^{-2}\nu/2)^{1-p}\Big(\int_{z-\frac{1}{4}K^{-2}\nu}^{z+\frac{1}{4}K^{-2}\nu} |u(x+\ell)-u(x)| dx\Big)^p \\
&= C(p)\nu^{1-p}\Big(\int_{z-\frac{1}{4}K^{-2}\nu}^{z+\frac{1}{4}K^{-2}\nu} \Big(\int_x^{x+\ell} -u'(y)dy\Big)dx\Big)^p \\
&\geq C(p)\nu^{1-p}\Big(\int_{z-\frac{1}{4}K^{-2}\nu}^{z+\frac{1}{4}K^{-2}\nu} \frac{1}{2}\ell K^{-1}\nu^{-1}\ dx\Big)^p = C(p)\nu^{1-p}\ell^p.
\end{aligned}$$

Case $\mathbf{p < 1}$. By Hölder's inequality we obtain that

$$\begin{aligned}
&\int_{S^1} |u(x+\ell)-u(x)|^p dx \geq \int_{S^1} \Big((u(x+\ell)-u(x))^+\Big)^p dx \\
&\geq \Big(\int_{S^1} \Big((u(x+\ell)-u(x))^+\Big)^2 dx\Big)^{p-1} \Big(\int_{S^1} (u(x+\ell)-u(x))^+ dx\Big)^{2-p}.
\end{aligned}$$

Using the upper estimate in (5.5) we get

$$\begin{aligned}
&\int_{S^1} |u(x+\ell)-u(x)|^p dx \\
&\geq \Big(\int_{S^1} \ell^2K^2 dx\Big)^{p-1} \Big(\int_{S^1} (u(x+\ell)-u(x))^+ dx\Big)^{2-p}.
\end{aligned}$$

Since $\int_{S^1} (u(\cdot+\ell)-u(\cdot)) = 0$, we obtain that

$$\begin{aligned}
&\int_{S^1} |u(x+\ell)-u(x)|^p dx \\
&\geq C(p)\ell^{2(p-1)}\Big(\frac{1}{2}\int_{S^1} |u(x+\ell)-u(x)| dx\Big)^{2-p} \geq C(p)\ell^p.
\end{aligned}$$

The last inequality follows from the case $p = 1$. □

The proof of the following lemma uses an argument from Aurell, Frisch, Lutsko & Vergassola (1992), which becomes quantitative if we restrict ourselves to the set O_K.

Lemma 5.8. *For $m \geq 0$ and $\ell \in J_2$,*

$$S_p(\ell) \overset{p}{\gtrsim} \begin{cases} \ell^p, & 0 \leq p \leq 1. \\ \ell, & p \geq 1. \end{cases}$$

Proof. In the same way as above, it suffices to prove that the inequalities hold uniformly in t for $t \in O_K$, with $S_p(\ell)$ replaced by

$$\int_{S^1} |u(x+\ell)-u(x)|^p dx,$$

and we can restrict ourselves to the case $p \geq 1$. Again, till the end of this proof, we assume that $t \in O_K$.

Defining z as in the proof of Lemma 5.7, we have

$$\int_{S^1} |u(x+\ell) - u(x)|^p dx \geq \int_{z-\frac{1}{2}\ell}^{z} \Big| \int_x^{x+\ell} u'^-(y)dy - \int_x^{x+\ell} u'^+(y)dy \Big|^p dx.$$

Since $\ell \geq C_1\nu = \frac{1}{4}K^{-2}\nu$, then by (5.12) for $x \in [z - \frac{1}{2}\ell, z]$ we get

$$\int_x^{x+\ell} u'^-(y)dy \geq \int_z^{z+\frac{1}{8}K^{-2}\nu} u'^-(y)dy \geq \frac{1}{16}K^{-3}.$$

.

On the other hand, since $\ell \leq C_2$, then by (5.5) and (5.10) we have

$$\int_x^{x+\ell} u'^+(y)dy \leq C_2 K = \frac{1}{20}K^{-3}.$$

Thus,

$$\int_{S^1} |u(x+\ell) - u(x)|^p dx \geq \frac{1}{2}\ell\left(\Big(\frac{1}{16} - \frac{1}{20}\Big)K^{-3}\right)^p \geq C(p)\ell. \ \Box$$

Summing up the results above we obtain the following theorem.

Theorem 5.9. *For $\ell \in J_1$,*

$$S_p(\ell) \overset{p}{\sim} \begin{cases} \ell^p, \ 0 \leq p \leq 1. \\ \ell^p \nu^{-(p-1)}, \ p \geq 1. \end{cases}$$

On the other hand, for $\ell \in J_2$,

$$S_p(\ell) \overset{p}{\sim} \begin{cases} \ell^p, \ 0 \leq p \leq 1. \\ \ell, \ p \geq 1. \end{cases}$$

The following result follows immediately from the definition (5.2).

Corollary 5.10. *For $\ell \in J_2$, the flatness satisfies $F(\ell) \sim \ell^{-1}$.*

It remains to prove that, as long as $|k|$ remains in a certain range, after layer-averaging, we have $\{|\hat{u}_k|^2\} \sim |k|^{-2}$. For this, we use a version of the Wiener-Khinchin theorem, stating that for any function $v \in L_2$ one has

$$|v(\cdot + y) - v(\cdot)|^2 = 4 \sum_{n \in \mathbb{Z}} \sin^2(\pi n y) |\hat{v}_n|^2. \tag{5.13}$$

Theorem 5.11. *For k such that $k^{-1} \in J_2$, we have $E(k) \sim k^{-2}$.*

Proof. We recall that by definition (5.3),

$$E(k) = \left\{ \frac{\sum_{|n| \in [M^{-1}k, Mk]} |\hat{u}_n|^2}{\sum_{|n| \in [M^{-1}k, Mk]} 1} \right\}.$$

Therefore proving the assertion of the theorem is the same as proving that

$$\sum_{|n|\in[M^{-1}k,Mk]} n^2\{|\hat{u}_n|^2\} \sim k. \tag{5.14}$$

The upper estimate holds without averaging over n such that $|n|\in[M^{-1}k,Mk]$. Indeed, by (5.1) we know that

$$\{|\hat{u}_n|^2\} \leq Cn^{-2},$$

which proves the upper bound. Also, this inequality implies that

$$\sum_{|n|<M^{-1}k} n^2\{|\hat{u}_n|^2\} \leq CM^{-1}k \tag{5.15}$$

and

$$\sum_{|n|>Mk} \{|\hat{u}_n|^2\} \leq CM^{-1}k^{-1}. \tag{5.16}$$

To prove the lower bound we note that

$$\begin{aligned}\sum_{|n|\leq Mk} n^2\{|\hat{u}_n|^2\} &\geq \frac{k^2}{\pi^2}\sum_{|n|\leq Mk}\sin^2(\pi nk^{-1})\{|\hat{u}_n|^2\}\\ &\geq \frac{k^2}{\pi^2}\Big(\sum_{n\in\mathbb{Z}}\sin^2(\pi nk^{-1})\{|\hat{u}_n|^2\} - \sum_{|n|>Mk}\{|\hat{u}_n|^2\}\Big).\end{aligned}$$

Using (5.13) and (5.16) we get

$$\begin{aligned}\sum_{|n|\leq Mk} n^2\{|\hat{u}_n|^2\} &\geq \frac{k^2}{4\pi^2}\Big(\{|u(\cdot+k^{-1})-u(\cdot)|^2\} - CM^{-1}k^{-1}\Big)\\ &\geq \frac{k^2}{4\pi^2}\Big(S_2(k^{-1}) - CM^{-1}k^{-1}\Big).\end{aligned}$$

Finally, using Theorem 5.9 we obtain that

$$\sum_{|n|\leq Mk} n^2\{|\hat{u}_n|^2\} \geq (C-CM^{-1})k.$$

Now we use (5.15) and we choose $M\geq 1$ large enough to obtain (5.14). □

6. Estimates for Small-Scale Quantities in the Inviscid Limit

It is a well-known fact (see for instance Kreiss & Lorenz (1989)) that as ν tends to 0, the solutions of (1.2) converge to *weak entropy solutions* of the inviscid equation $u_t + f'(u)u_x = 0$, for fixed t. The convergence takes place for almost every x, and therefore also in L_1, since solutions are uniformly bounded for all ν.

Those solutions, denoted u^0, inherit all previously proved properties which hold uniformly for small enough ν for solutions of (1.2).

To begin with, we define the non-empty and non-intersecting intervals

$$J_2 = (0,\ C_2];\ J_3 = (C_2,\ 1],$$

which now correspond to the inertial range and the energy range, respectively: the inviscid Burgers equation does not have a dissipation range, since formally there is no damping term. The constant C_2 is the same as above.

Then we define S_p^0, F^0 and E^0 for solutions $u^0(t,x)$ in the same way as the previously considered quantities S_p, F and E for solutions of the viscous equation. By the dominated convergence theorem, we obtain the following results:

THEOREM 6.1. *For* $\ell \in J_2$, $S_p^0(\ell) \overset{p}{\sim} \begin{cases} \ell^p, \ 0 \leq p \leq 1. \\ \ell, \ p \geq 1. \end{cases}$

COROLLARY 6.2. *For* $\ell \in J_2$, *the flatness satisfies* $F^0(\ell) \sim \ell^{-1}$.

THEOREM 6.3. *For* k *such that* $k^{-1} \in J_2$, *we have* $E^0(k) \sim k^{-2}$.

I would like to thank my Ph.D. advisor S.Kuksin, who originally formulated the problem. I am also very grateful to A.Biryuk, in particular for bringing to my attention some articles of S.Kruzhkov, and to J.Bec, U.Frisch, and K.Khanin for many helpful discussions. Finally, I would like to thank all of the staff and faculty at the CMLS in Ecole Polytechnique for advice, support, and discussions.

REFERENCES

AURELL, E., FRISCH, U., LUTSKO, J. & VERGASSOLA, M. 1992 On the multifractal properties of the energy dissipation derived from turbulence data. *Journal of Fluid Mechanics* **238**, 467–486.

BEC, J. & FRISCH, U. 2001 Burgulence. In *Les Houches 2000: New Trends in Turbulence* (ed. M. Lesieur, A.Yaglom & F. David), pp. 341–383. Springer EDP-Sciences.

BEC, J. & KHANIN, K. 2007 Burgers turbulence. *Physics Reports* **447**, 1–66.

BIRYUK, A. 2001 Spectral properties of solutions of the Burgers equation with small dissipation. *Functional Analysis and its Applications* **35:1**, 1–12.

BORITCHEV, A. 2012*a* Estimates for solutions of a low-viscosity kick-forced generalised Burgers equation. *To appear in Proceedings of the Royal Society of Edinburgh A* ArXiv:1107.4866v1.

BORITCHEV, A. 2012*b* Sharp estimates for turbulence in white-forced generalised Burgers equation. ArXiv:1201.5567.

BURGERS, J. M. 1974 *The nonlinear diffusion equation: asymptotic solutions and statistical problems*, *University of Maryland, Institute for Fluid Dynamics and Applied Mathematics Lecture Series*, vol. 52. Reidel.

CHORIN, A. 1975 *Lectures on turbulence theory*, *Mathematics Lecture Series*, vol. 5. Publish or Perish.

DOERING, C. & GIBBON, J. D. 1995 *Applied analysis of the Navier-Stokes equations*. Cambridge University Press.

E, WEINAN, KHANIN, K., MAZEL, A. & SINAI, YA. 1997 Probability distribution functions for the random forced Burgers equation. *Physical Review Letters* (78:10), 1904–1907.

EVANS, L. 2008 *Partial differential equations*, *AMS Graduate Studies in Mathematics*, vol. 19.

FLORIN, V. 1948 Some of the simplest nonlinear problems arising in the consolidation of wet soil. *Izvestiya Akademii Nauk SSSR Otdel Technicheskih Nauk* (9), 1389–1402.

FOURNIER, J. D. & FRISCH, U. 1983 L'équation de Burgers déterministe et stastistique. *Journal de Mécanique Théorique et Appliquée* (2), 699–750.

FRISCH, U. 1995 *Turbulence: the legacy of A.N. Kolmogorov*. Cambridge University Press.

KIDA, S. 1979 Asymptotic properties of Burgers turbulence. *Journal of Fluid Mechanics* (93:2), 337–377.

KOLMOGOROV, A. 1941*a* Dissipation of energy in locally isotropic turbulence. *Doklady Akademii Nauk SSSR* (32), 16–18, reprinted in Proceedings of the Royal Society of London A 434 (1991), 15-17.

KOLMOGOROV, A. 1941*b* On degeneration (decay) of isotropic turbulence in an incompressible viscous liquid. *Doklady Akademii Nauk SSSR* (31), 538–540.

KOLMOGOROV, A. 1941*c* The local structure of turbulence in incompressible viscous fluid for very large Reynolds number. *Doklady Akademii Nauk SSSR* (30), 9–13, reprinted in Proceedings of the Royal Society of London A 434 (1991), 9-13.

KRAICHNAN, R. H. 1968 Lagrangian-history statistical theory for Burgers' equation. *Physics of Fluids* (11:2), 265–277.

KREISS, H.-O. & LORENZ, J. 1989 *Initial-boundary value problems and the Navier-Stokes equations*, *Pure and Applied Mathematics*, vol. 136. Academic Press.

KRUZHKOV, S. N. 1964 The Cauchy Problem in the large for nonlinear equations and for certain quasilinear systems of the first-order with several variables. *Soviet Math. Doklady* (5), 493–496.

KUKSIN, S. 1997 On turbulence in nonlinear Schrödinger equations. *Geometric and Functional Analysis* (7), 783–822.

KUKSIN, S. 1999 Spectral properties of solutions for nonlinear PDEs in the turbulent regime. *Geometric and Functional Analysis* (9), 141–184.